

References

1. Wu, M. Y., and Hill, C. S. (2009) TGF- β superfamily signaling in embryonic development and homeostasis. *Dev Cell* **16**, 329-343
2. Weiss, A., and Attisano, L. (2013) The TGF β superfamily signaling pathway. *Wiley Interdiscip Rev Dev Biol* **2**, 47-63
3. Gray, A. M., and Mason, A. J. (1990) Requirement for activin A and transforming growth factor- β 1 pro-regions in homodimer assembly. *Science* **247**, 1328-1330
4. Sha, X., Yang, L., and Gentry, L. E. (1991) Identification and analysis of discrete functional domains in the pro region of pre-pro-transforming growth factor beta 1. *J Cell Biol* **114**, 827-839
5. Neugebauer, J. M., Kwon, S., Kim, H. S., Donley, N., Tilak, A., Sopory, S., and Christian, J. L. (2015) The prodomain of BMP4 is necessary and sufficient to generate stable BMP4/7 heterodimers with enhanced bioactivity in vivo. *Proc Natl Acad Sci U S A* **112**, E2307-2316
6. Miyazono, K., Hellman, U., Wernstedt, C., and Heldin, C. H. (1988) Latent high molecular weight complex of transforming growth factor beta 1. Purification from human platelets and structural characterization. *J Biol Chem* **263**, 6407-6415
7. Saharinen, J., and Keski-Oja, J. (2000) Specific sequence motif of 8-Cys repeats of TGF- β binding proteins, LTBP, creates a hydrophobic interaction surface for binding of small latent TGF- β . *Mol Biol Cell* **11**, 2691-2704
8. Gregory, K. E., Ono, R. N., Charbonneau, N. L., Kuo, C. L., Keene, D. R., Bachinger, H. P., and Sakai, L. Y. (2005) The prodomain of BMP-7 targets the BMP-7 complex to the extracellular matrix. *J Biol Chem* **280**, 27970-27980
9. Sengle, G., Charbonneau, N. L., Ono, R. N., Sasaki, T., Alvarez, J., Keene, D. R., Bachinger, H. P., and Sakai, L. Y. (2008) Targeting of bone morphogenetic protein growth factor complexes to fibrillin. *J Biol Chem* **283**, 13874-13888
10. Li, S., Shimono, C., Norioka, N., Nakano, I., Okubo, T., Yagi, Y., Hayashi, M., Sato, Y., Fujisaki, H., Hattori, S., Sugiura, N., Kimata, K., and Sekiguchi, K. (2010) Activin A binds to perlecan through its pro-region that has heparin/heparan sulfate binding activity. *J Biol Chem* **285**, 36645-36655
11. Anderson, S. B., Goldberg, A. L., and Whitman, M. (2008) Identification of a novel pool of extracellular pro-myostatin in skeletal muscle. *J Biol Chem* **283**, 7027-7035
12. Bauskin, A. R., Brown, D. A., Junankar, S., Rasiah, K. K., Eggleton, S., Hunter, M., Liu, T., Smith, D., Kuffner, T., Pankhurst, G. J., Johnen, H., Russell, P. J., Barret, W., Stricker, P. D., Grygiel, J. J., Kench, J. G., Henshall, S. M., Sutherland, R. L., and Breit, S. N. (2005) The propeptide mediates formation of stromal stores of PROMIC-1: role in determining prostate cancer outcome. *Cancer Res* **65**, 2330-2336

13. Stockis, J., Colau, D., Coulie, P. G., and Lucas, S. (2009) Membrane protein GARP is a receptor for latent TGF- β on the surface of activated human Treg. *Eur J Immunol* **39**, 3315-3322
14. Tran, D. Q., Andersson, J., Wang, R., Ramsey, H., Unutmaz, D., and Shevach, E. M. (2009) GARP (LRRC32) is essential for the surface expression of latent TGF- β on platelets and activated FOXP3+ regulatory T cells. *Proc Natl Acad Sci U S A* **106**, 13445-13450
15. Harrison, C. A., Al-Musawi, S. L., and Walton, K. L. (2011) Prodomains regulate the synthesis, extracellular localisation and activity of TGF- β superfamily ligands. *Growth Factors* **29**, 174-186
16. Wang, R., Zhu, J., Dong, X., Shi, M., Lu, C., and Springer, T. A. (2012) GARP regulates the bioavailability and activation of TGF- β . *Mol. Biol. Cell* **23**, 1129-1139
17. Qin, Y., Garrison, B. S., Ma, W., Wang, R., Jiang, A., Li, J., Mistry, M., Bronson, R. T., Santoro, D., Franco, C., Robinton, D. A., Stevens, B., Rossi, D. J., Lu, C., and Springer, T. A. (2018) A milieu molecule for TGF- β required for microglia function in the nervous system. *Cell* **174**, 156-171 e116
18. Cui, Y., Hackenmiller, R., Berg, L., Jean, F., Nakayama, T., Thomas, G., and Christian, J. L. (2001) The activity and signaling range of mature BMP-4 is regulated by sequential cleavage at two sites within the prodomain of the precursor. *Genes Dev* **15**, 2797-2802
19. Degnin, C., Jean, F., Thomas, G., and Christian, J. L. (2004) Cleavages within the prodomain direct intracellular trafficking and degradation of mature bone morphogenetic protein-4. *Mol Biol Cell* **15**, 5012-5020
20. Sopory, S., Kwon, S., Wehrli, M., and Christian, J. L. (2010) Regulation of Dpp activity by tissue-specific cleavage of an upstream site within the prodomain. *Dev Biol* **346**, 102-112
21. Akiyama, T., Marques, G., and Wharton, K. A. (2012) A large bioactive BMP ligand with distinct signaling properties is produced by alternative proconvertase processing. *Sci Signal* **5**, ra28
22. Kunnappu, J., Tauscher, P. M., Tiusanen, N., Nguyen, M., Loytynoja, A., Arora, K., and Shimmi, O. (2014) Cleavage of the Drosophila screw prodomain is critical for a dynamic BMP morphogen gradient in embryogenesis. *Dev Biol* **389**, 149-159
23. Mi, L. Z., Brown, C. T., Gao, Y., Tian, Y., Le, V. Q., Walz, T., and Springer, T. A. (2015) Structure of bone morphogenetic protein 9 procomplex. *Proc Natl Acad Sci U S A* **112**, 3710-3715
24. Anderson, E. N., and Wharton, K. A. (2017) Alternative cleavage of the bone morphogenetic protein (BMP), Gbb, produces ligands with distinct developmental functions and receptor preferences. *J Biol Chem* **292**, 19160-19178

25. Simpson, C. M., Stanton, P. G., Walton, K. L., Chan, K. L., Ritter, L. J., Gilchrist, R. B., and Harrison, C. A. (2012) Activation of latent human GDF9 by a single residue change (Gly 391 Arg) in the mature domain. *Endocrinology* **153**, 1301-1310
26. Gentry, L. E., Webb, N. R., Lim, G. J., Brunner, A. M., Ranchalis, J. E., Twardzik, D. R., Lioubin, M. N., Marquardt, H., and Purchio, A. F. (1987) Type 1 transforming growth factor beta: amplified expression and secretion of mature and precursor polypeptides in Chinese hamster ovary cells. *Mol Cell Biol* **7**, 3418-3427
27. Wakefield, L. M., Smith, D. M., Flanders, K. C., and Sporn, M. B. (1988) Latent transforming growth factor-beta from human platelets. A high molecular weight complex containing precursor sequences. *J Biol Chem* **263**, 7646-7654
28. Khalil, N. (1999) TGF-beta: from latent to active. *Microbes Infect* **1**, 1255-1263
29. Lee, S. J., and McPherron, A. C. (2001) Regulation of myostatin activity and muscle growth. *Proc Natl Acad Sci U S A* **98**, 9306-9311
30. Thies, R. S., Chen, T., Davies, M. V., Tomkinson, K. N., Pearson, A. A., Shakey, Q. A., and Wolfman, N. M. (2001) GDF-8 propeptide binds to GDF-8 and antagonizes biological activity by inhibiting GDF-8 receptor binding. *Growth Factors* **18**, 251-259
31. Hill, J. J., Davies, M. V., Pearson, A. A., Wang, J. H., Hewick, R. M., Wolfman, N. M., and Qiu, Y. (2002) The myostatin propeptide and the follistatin-related gene are inhibitory binding proteins of myostatin in normal serum. *J Biol Chem* **277**, 40735-40741
32. Zimmers, T. A., Davies, M. V., Koniaris, L. G., Haynes, P., Esquela, A. F., Tomkinson, K. N., McPherron, A. C., Wolfman, N. M., and Lee, S. J. (2002) Induction of cachexia in mice by systemically administered myostatin. *Science* **296**, 1486-1488
33. Ge, G., Hopkins, D. R., Ho, W. B., and Greenspan, D. S. (2005) GDF11 forms a bone morphogenetic protein 1-activated latent complex that can modulate nerve growth factor-induced differentiation of PC12 cells. *Mol Cell Biol* **25**, 5846-5858
34. Sengle, G., Ono, R. N., Sasaki, T., and Sakai, L. Y. (2011) Prodomains of transforming growth factor β (TGF β) superfamily members specify different functions: extracellular matrix interactions and growth factor bioavailability. *J Biol Chem* **286**, 5087-5099
35. Munger, J. S., Huang, X., Kawakatsu, H., Griffiths, M. J. D., Dalton, S. L., Wu, J., Pittet, J.-F., Kaminski, N., Garat, C., Matthay, M. A., Rifkin, D. B., and Sheppard, D. (1999) The integrin $\alpha\beta 6$ binds and activates latent TGF $\beta 1$: A mechanism for regulating pulmonary inflammation and fibrosis. *Cell* **96**, 319-328
36. Mu, D., Cambier, S., Fjellbirkeland, L., Baron, J. L., Munger, J. S., Kawakatsu, H., Sheppard, D., Broaddus, V. C., and Nishimura, S. L. (2002) The integrin $\alpha\beta 8$ mediates epithelial homeostasis through MT1-MMP-dependent activation of TGF- $\beta 1$. *J. Cell Biol.* **157**, 493-507
37. Dong, X., Zhao, B., Iacob, R. E., Zhu, J., Koksai, A. C., Lu, C., Engen, J. R., and Springer, T. A. (2017) Force interacts with macromolecular structure in activation of TGF- β . *Nature* **542**, 55-59

38. Wolfman, N. M., McPherron, A. C., Pappano, W. N., Davies, M. V., Song, K., Tomkinson, K. N., Wright, J. F., Zhao, L., Sebald, S. M., Greenspan, D. S., and Lee, S. J. (2003) Activation of latent myostatin by the BMP-1/tolloid family of metalloproteinases. *Proc Natl Acad Sci U S A* **100**, 15842-15846
39. Sampath, T. K., and Reddi, A. H. (1981) Dissociative extraction and reconstitution of extracellular matrix components involved in local bone differentiation. *Proc Natl Acad Sci U S A* **78**, 7599-7603
40. Wang, E. A., Rosen, V., Cordes, P., Hewick, R. M., Kriz, M. J., Luxenberg, D. P., Sibley, B. S., and Wozney, J. M. (1988) Purification and characterization of other distinct bone-inducing factors. *Proc Natl Acad Sci U S A* **85**, 9484-9488
41. Luyten, F. P., Cunningham, N. S., Ma, S., Muthukumaran, N., Hammonds, R. G., Nevins, W. B., Woods, W. I., and Reddi, A. H. (1989) Purification and partial amino acid sequence of osteogenin, a protein initiating bone differentiation. *J Biol Chem* **264**, 13377-13380
42. Celeste, A. J., Iannazzi, J. A., Taylor, R. C., Hewick, R. M., Rosen, V., Wang, E. A., and Wozney, J. M. (1990) Identification of transforming growth factor beta family members present in bone-inductive protein purified from bovine bone. *Proc Natl Acad Sci U S A* **87**, 9843-9847
43. Sampath, T. K., Coughlin, J. E., Whetstone, R. M., Banach, D., Corbett, C., Ridge, R. J., Ozkaynak, E., Oppermann, H., and Rueger, D. C. (1990) Bovine osteogenic protein is composed of dimers of OP-1 and BMP-2A, two members of the transforming growth factor-beta superfamily. *J Biol Chem.* **265**, 13198-13205
44. Wohl, A. P., Troilo, H., Collins, R. F., Baldock, C., and Sengle, G. (2016) Extracellular Regulation of Bone Morphogenetic Protein Activity by the Microfibril Component Fibrillin-1. *J Biol Chem* **291**, 12732-12746
45. Le, V. Q., Iacob, R. E., Tian, Y., McConaughy, W., Jackson, J., Su, Y., Zhao, B., Engen, J. R., Pirruccello-Straub, M., and Springer, T. A. (2018) Tolloid cleavage activates latent GDF8 by priming the pro-complex for dissociation. *EMBO J* **37**, 384-397
46. Cotton, T. R., Fischer, G., Wang, X., McCoy, J. C., Czepnik, M., Thompson, T. B., and Hyvonen, M. (2018) Structure of the human myostatin precursor and determinants of growth factor latency. *EMBO J* **37**, 367-383
47. Shi, M., Zhu, J., Wang, R., Chen, X., Mi, L. Z., Walz, T., and Springer, T. A. (2011) Latent TGF- β structure and activation. *Nature* **474**, 343-349
48. Zhao, B., Xu, S., Dong, X., Lu, C., and Springer, T. A. (2018) Prodomain-growth factor swapping in the structure of pro-TGF- β 1. *J Biol Chem* **293**, 1579-1589
49. Wang, X., Fischer, G., and Hyvonen, M. (2016) Structure and activation of pro-activin A. *Nat Commun* **7**, 12052

50. Fang, J., Nevin, P., Kairys, V., Venclovas, C., Engen, J. R., and Beuning, P. J. (2014) Conformational analysis of processivity clamps in solution demonstrates that tertiary structure does not correlate with protein dynamics. *Structure* **22**, 572-581
51. Wales, T. E., Poe, J. A., Emert-Sedlak, L., Morgan, C. R., Smithgall, T. E., and Engen, J. R. (2016) Hydrogen exchange mass spectrometry of related proteins with divergent sequences: A comparative study of HIV-1 nef allelic variants. *J Am Soc Mass Spectrom* **27**, 1048-1061
52. Harrison, R. A., Lu, J., Carrasco, M., Hunter, J., Manandhar, A., Gondi, S., Westover, K. D., and Engen, J. R. (2016) Structural dynamics in Ras and related proteins upon nucleotide switching. *J Mol Biol* **428**, 4723-4735
53. Wang, J., Su, Y., Iacob, R. E., Engen, J. R., and Springer, T. A. (2019) General structural features that regulate integrin affinity revealed by atypical $\alpha V\beta 8$. *Nat Commun* **10**, 5481
54. Kochert, B. A., Fleischhacker, A. S., Wales, T. E., Becker, D. F., Engen, J. R., and Ragsdale, S. W. (2019) Dynamic and structural differences between heme oxygenase-1 and -2 are due to differences in their C-terminal regions. *J Biol Chem* **294**, 8259-8272
55. Hinck, A. P., Mueller, T. D., and Springer, T. A. (2016) Structural Biology and Evolution of the TGF- β Family. *Cold Spring Harb Perspect Biol* **8**
56. Lienart, S., Merceron, R., Vanderaa, C., Lambert, F., Colau, D., Stockis, J., van der Woning, B., De Haard, H., Saunders, M., Coulie, P. G., Savvides, S. N., and Lucas, S. (2018) Structural basis of latent TGF- $\beta 1$ presentation and activation by GARP on human regulatory T cells. *Science* **362**, 952-956
57. Jiang, M. S., Liang, L. F., Wang, S., Ratovitski, T., Holmstrom, J., Barker, C., and Stotish, R. (2004) Characterization and identification of the inhibitory domain of GDF-8 propeptide. *Biochem Biophys Res Commun* **315**, 525-531
58. Ohsawa, Y., Takayama, K., Nishimatsu, S., Okada, T., Fujino, M., Fukai, Y., Murakami, T., Hagiwara, H., Itoh, F., Tsuchida, K., Hayashi, Y., and Sunada, Y. (2015) The Inhibitory Core of the Myostatin Prodomain: Its Interaction with Both Type I and II Membrane Receptors, and Potential to Treat Muscle Atrophy. *PLoS One* **10**, e0133713
59. Takayama, K., Noguchi, Y., Aoki, S., Takayama, S., Yoshida, M., Asari, T., Yakushiji, F., Nishimatsu, S., Ohsawa, Y., Itoh, F., Negishi, Y., Sunada, Y., and Hayashi, Y. (2015) Identification of the minimum peptide from mouse myostatin prodomain for human myostatin inhibition. *J Med Chem* **58**, 1544-1549
60. Chen, J. L., Walton, K. L., Al-Musawi, S. L., Kelly, E. K., Qian, H., La, M., Lu, L., Lovrecz, G., Ziemann, M., Lazarus, R., El-Osta, A., Gregorevic, P., and Harrison, C. A. (2015) Development of novel activin-targeted therapeutics. *Mol Ther* **23**, 434-444
61. Walton, K. L., Makanji, Y., Chen, J., Wilce, M. C., Chan, K. L., Robertson, D. M., and Harrison, C. A. (2010) Two distinct regions of latency-associated peptide coordinate stability of the latent transforming growth factor-beta1 complex. *J Biol Chem* **285**, 17029-17037

62. Walker, R. G., McCoy, J. C., Czepnik, M., Mills, M. J., Hagg, A., Walton, K. L., Cotton, T. R., Hyvonen, M., Lee, R. T., Gregorevic, P., Harrison, C. A., and Thompson, T. B. (2018) Molecular characterization of latent GDF8 reveals mechanisms of activation. *Proc Natl Acad Sci U S A* **115**, E866-E875
63. Rees, I., Langley, E., Chiu, W., and Ludtke, S. J. (2013) EMEN2: an object oriented database and electronic lab notebook. *Microsc Microanal* **19**, 1-10
64. Chen, X., Xie, C., Nishida, N., Li, Z., Walz, T., and Springer, T. A. (2010) Requirement of open headpiece conformation for activation of leukocyte integrin alphaXbeta2. *Proc Natl Acad Sci U S A* **107**, 14727-14732
65. Frank, J., Radermacher, M., Penczek, P., Zhu, J., Li, Y., Ladjadj, M., and Leith, A. (1996) SPIDER and WEB: processing and visualization of images in 3D electron microscopy and related fields. *J. Struct. Biol.* **116**, 190-199
66. Mi, L. Z., Lu, C., Nishida, N., Walz, T., and Springer, T. A. (2011) Simultaneous visualization of the extracellular and cytoplasmic domains of the epidermal growth factor receptor. *Nat. Struct. Biol.* **18**, 984-989
67. Iacob, R. E. B.-A., G. M.; Makowski, L.; Engen, J. R.; Berkowitz, S. A.; Houde, D. (2013) Investigating monoclonal antibody aggregation using a combination of H/DX-MS and other biophysical measurements. *J Pharm Sci* **102**, 4315-4329
68. Masson, G. R., Burke, J. E., Ahn, N. G., Anand, G. S., Borchers, C., Brier, S., Bou-Assaf, G. M., Engen, J. R., Englander, S. W., Faber, J., Garlish, R., Griffin, P. R., Gross, M. L., Guttman, M., Hamuro, Y., Heck, A. J. R., Houde, D., Iacob, R. E., Jorgensen, T. J. D., Kaltashov, I. A., Klinman, J. P., Konermann, L., Man, P., Mayne, L., Pascal, B. D., Reichmann, D., Skehel, M., Snijder, J., Strutzenberg, T. S., Underbakke, E. S., Wagner, C., Wales, T. E., Walters, B. T., Weis, D. D., Wilson, D. J., Wintrode, P. L., Zhang, Z., Zheng, J., Schriemer, D. C., and Rand, K. D. (2019) Recommendations for performing, interpreting and reporting hydrogen deuterium exchange mass spectrometry (HDX-MS) experiments. *Nat Methods* **16**, 595-602
69. Wales, T. E. F., K. E.; Gerhardt, G. C.; Engen, J. R. (2008) High-speed and high-resolution UPLC separation at zero degrees Celsius. *Anal Chem* **80**, 6815-6820
70. Wales, T. E., and Engen, J. R. (2006) Hydrogen exchange mass spectrometry for the analysis of protein dynamics. *Mass Spectrometry Reviews* **25**, 158-170
71. Perez-Riverol, Y., Csordas, A., Bai, J., Bernal-Llinares, M., Hewapathirana, S., Kundu, D. J., Inuganti, A., Griss, J., Mayer, G., Eisenacher, M., Perez, E., Uszkoreit, J., Pfeuffer, J., Sachsenberg, T., Yilmaz, S., Tiwary, S., Cox, J., Audain, E., Walzer, M., Jarnuczak, A. F., Ternent, T., Brazma, A., and Vizcaino, J. A. (2019) The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Res* **47**, D442-D450