



RESEARCH ARTICLE

Dendritic cell-expressed common gamma-chain recruits IL-15 for trans-presentation at the murine immunological synapse

[version 1; referees: 1 approved, 1 approved with reservations]

Chiara Beilin ^{1*}, Kaushik Choudhuri^{2*}, Gerben Bouma ¹, Dessislava Malinova ¹, Jaime Llodra³, David L. Stokes³, Motumu Shimaoka⁴, Timothy A. Springer ³, Michael L. Dustin ^{2,5}, Adrian J. Thrasher^{1,6}, Siobhan O. Burns ^{6,7}

¹Molecular Immunology Unit, Institute of Child Health, University College London, London, WC1N 1EH, UK

²Program in Molecular Pathogenesis, Skirball Institute of Biomolecular Medicine, New York University, New York, NY, 10016, USA

³Program in Structural Biology, Skirball Institute of Biomolecular Medicine, New York University, New York, NY, 10016, USA

⁴Immune Disease Institute, Children's Hospital Boston, Boston, MA, 02115, USA

⁵Kennedy Institute of Rheumatology, Nuffield Department of Orthopedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Headington, OX3 7FY, UK

⁶Great Ormond Street Hospital for Children NHS Foundation Trust, London, WC1N 3JH, UK

⁷University College London Institute of Immunity and Transplantation, Department of Immunology, Royal Free London NHS Foundation Trust, London, NW3 2PF, UK

* Equal contributors

v1 First published: 17 Jul 2018, 3:84 (doi: [10.12688/wellcomeopenres.14493.1](https://doi.org/10.12688/wellcomeopenres.14493.1))
Latest published: 17 Oct 2018, 3:84 (doi: [10.12688/wellcomeopenres.14493.2](https://doi.org/10.12688/wellcomeopenres.14493.2))

Abstract

Background: Mutations of the common cytokine receptor gamma chain (γ_c) cause Severe Combined Immunodeficiency characterized by absent T and NK cell development. Although stem cell therapy restores these lineages, residual immune defects are observed that may result from selective persistence of γ_c -deficiency in myeloid lineages. However, little is known about the contribution of myeloid-expressed γ_c to protective immune responses. Here we examine the importance of γ_c for myeloid dendritic cell (DC) function.

Methods: We utilize a combination of *in vitro* DC/T-cell co-culture assays and a novel lipid bilayer system mimicking the T cell surface to delineate the role of DC-expressed γ_c during DC/T-cell interaction.

Results: We observed that γ_c in DC was recruited to the contact interface following MHCII ligation, and promoted IL-15R α colocalization with engaged MHCII. Unexpectedly, trans-presentation of IL-15 was required for optimal CD4+T cell activation by DC and depended on DC γ_c expression. Neither recruitment of IL-15R α nor IL-15 trans-signaling at the DC immune synapse (IS), required γ_c signaling in DC, suggesting that γ_c facilitates IL-15 transpresentation through induced intermolecular *cis* associations or cytoskeletal reorganization following MHCII ligation.

Conclusions: These findings show that DC-expressed γ_c is required for effective antigen-induced CD4+ T cell activation. We reveal a novel mechanism for recruitment of DC IL-15/IL-15R α complexes to the IS, leading to CD4+ T cell costimulation through localized IL-15 transpresentation that is coordinated with antigen-recognition.

Open Peer Review

Referee Status:

Invited Referees		
	1	2
version 2 published 17 Oct 2018		
version 1 published 17 Jul 2018	 report	 report

- Matthew Collin** , Newcastle University, UK
Newcastle Hospitals NHS Foundation Trust, UK
- Jung-Hyun Park** , NIH, USA
Tae-Hyoun Kim, National Institutes of Health, USA

Keywords

interleukins, immunological synapse, immunodeficiency, trans-presentation, dendritic cells, lymphocytes,

Discuss this article

Comments (0)

Corresponding authors: Michael L. Dustin (mikeroscopedustin@gmail.com), Siobhan O. Burns (siobhan.burns@ucl.ac.uk)

Author roles: **Beilin C:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation; **Choudhuri K:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Supervision, Writing – Original Draft Preparation; **Bouma G:** Formal Analysis, Investigation, Methodology, Writing – Review & Editing; **Malinova D:** Formal Analysis, Investigation, Methodology, Writing – Review & Editing; **Llodra J:** Formal Analysis, Investigation, Methodology, Writing – Review & Editing; **Stokes DL:** Funding Acquisition, Supervision, Writing – Review & Editing; **Shimaoka M:** Resources, Writing – Review & Editing; **Springer TA:** Funding Acquisition, Resources, Supervision, Writing – Review & Editing; **Dustin ML:** Conceptualization, Project Administration, Supervision, Writing – Review & Editing; **Thrasher AJ:** Funding Acquisition, Supervision, Writing – Review & Editing; **Burns SO:** Conceptualization, Formal Analysis, Funding Acquisition, Project Administration, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome Trust [090233, GB, AJT] and [100262Z, MLD]; the Child Health Research Appeal Trust and a Bogue Research Fellowship (CB); the Primary Immunodeficiency Association and the ICH Biomedical Research Centre (SOB); The European Union (GB); The Cancer Research Institute (KC); and US National Institutes of Health [grant R37 AI43542, MLD] and [R01 CA31798, TAS].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Beilin C *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Beilin C, Choudhuri K, Bouma G *et al.* **Dendritic cell-expressed common gamma-chain recruits IL-15 for trans-presentation at the murine immunological synapse [version 1; referees: 1 approved, 1 approved with reservations]** Wellcome Open Research 2018, 3:84 (doi: [10.12688/wellcomeopenres.14493.1](https://doi.org/10.12688/wellcomeopenres.14493.1))

First published: 17 Jul 2018, 3:84 (doi: [10.12688/wellcomeopenres.14493.1](https://doi.org/10.12688/wellcomeopenres.14493.1))

Introduction

Severe Combined Immunodeficiency (SCID) caused by deficiency of the common cytokine receptor gamma chain (γ_c) is characterized by defective T and NK cell development, resulting in life-threatening infections. Although the condition can be cured by bone marrow transplantation (BMT) or gene therapy, several long-term complications are seen; in particular a high incidence of severe cutaneous human papilloma virus (HPV) infection that suggests residual defects of immunity^{1–3}. HPV susceptibility is not predicted by transplantation conditions or subsequent immune reconstitution, but is curiously restricted to SCID resulting from mutations in γ_c or its signaling mediator Janus-associated kinase 3 (JAK3) and therefore appears to be related to the original genetic mutation.

HPV infections are limited to the epidermis suggesting persistent defects in the skin compartment, which could relate to keratinocytes or hematopoietic-derived immune cells. As many SCID patients receive BMT without any chemotherapy conditioning, B cell and myeloid lineages remain of host origin and therefore γ_c -deficient in the majority of cases^{1,2}. This includes antigen-presenting dendritic cells (DC) derived from bone marrow, such as dermal (migratory) DC⁴ and those that self-renew in tissues, such as epidermal Langerhans cells (LC). Although the mechanisms are poorly understood, LC and dermal DC are predicted to be important for regression of cutaneous HPV lesions through their role as potent skin antigen presenting cells for priming adaptive immune responses^{5,6}. It is thought that CD4⁺ T cells also play a central role in anti-HPV immunity, as their presence at sites of HPV infection are predictive of clearance, while susceptibility to HPV infection is dramatically increased by CD4⁺ T cell immunodeficiency^{7–11}. Since γ_c -deficiency in T cells is effectively corrected in SCID patients who have undergone BMT, we speculated that γ_c -deficient residual DC might be defective in priming antigen-specific CD4⁺ T cells in these patients, and hence might contribute to the observed impaired immunity to infection.

As *ex vivo* isolation of primary LC and dermal DC populations in large numbers is technically challenging, we modeled DC γ_c -deficiency using monocyte-derived DC generated from the bone marrow of γ_c -deficient mice. While DC subsets differ in specific functions that likely relate to the particular requirements of their tissue environments, all myeloid-derived DC populations share prototypical features, including antigen uptake, presentation and T cell priming⁴. DC/LC normally express several γ_c -containing cytokine receptors: specifically IL-2R, IL-4R, IL-15R and IL-21R that, upon binding their respective cytokines, regulate DC functions such as activation and cytokine release¹². In addition, DC-expressed IL-15R (and possibly IL-2R) regulates the function of other immune cells through the unusual mechanism of cytokine transpresentation that requires direct intercellular interaction^{13–15}. In particular, transpresentation of IL-15 by DC is required for NK cell and memory CD8⁺ T cell activation and homeostasis^{13,16}. Although several studies have shown that IL-15 enhances CD4⁺ T cell proliferation and is required for CD4⁺ memory homeostasis^{17–22} the importance of transpresentation for IL-15-dependent T-cell functions has not been clear until recently when effector CD4⁺ T-cell differentiation

was shown to rely on transpresented rather than soluble IL-15²³. To date, this has not been further detailed at a mechanistic level and a role for DC-mediated IL-15 transpresentation in CD4⁺ T-cell activation has not been documented.

In this study, we investigate the role of γ_c in DC function and identify a defect in the ability of γ_c -deficient DC to prime naïve CD4⁺ T cells. Using a novel supported planar bilayer system that mimics key molecular features of the T cell surface, we demonstrate that, independent of its signaling function, DC-expressed γ_c localises to the DC:T-cell contact interface following MHCII ligation and results in recruitment and colocalization of IL-15R α with MHCII. We show that γ_c -deficiency in DC critically impairs IL-15R α recruitment and IL-15 transpresentation to naïve CD4⁺ T cells at the immunological synapse, resulting in incomplete T cell activation. In light of these findings, we suggest a novel model for IL-15 transpresentation in which the DC-IS regulates co-stimulation of CD4⁺ T-cells during antigen-dependent priming.

Methods

Animals

Mice were C57BL/6 wild-type and OTII transgenic (OVA_{323–339} peptide (pOVA)/I-A^b-specific CD4⁺ T cells) (Charles River, Kent, UK). γ_c /Rag2^{-/-} mice (C57BL/6) were kindly provided by Dr. Colucci (Babraham Institute, Cambridge, UK). Male and female mice were housed in individually ventilated cages, up to 6 mice per cage with bedding changed twice weekly and sacrificed by exposure to a rising concentration of CO₂ at 8–12 weeks of age weighing approximately 25–30g. Bone marrow was extracted from tibia and femur bones. Work in mice was performed in an ethical manner according to UK Home Office regulations under project licence number PPL 70/7329.

Cloning procedures

The lentiviral construct encoding γ_c^{WT} -GFP fusion protein (pLV-CMVEI.hIL2RG-SceI-EGFPds) was kindly provided by Nadine Dannemann, Toni Cathomen Lab, Hannover Medical School. A truncated $\gamma_c^{\Delta C}$ -GFP was created by introduction of AgeI site by PCR with following primers: forward primer (GAAGACA CCGACTCTAGAGCCACCATGTTG), reverse primer (CAACCGGTGGGCATCGTCCGTTCCAG). The PCR product was digested with XbaI and AgeI and religated into the original vector to create a $\gamma_c^{\Delta C}$ -GFP fusion lacking 77 amino acids at the C terminus.

Cell isolation and culture

Bone marrow-derived DC (BMDC) were generated, LPS-matured and OVA-pulsed as previously described²⁴. BMDC were blocked for 30 min at 37°C with anti-IL-15R α (AF551) or isotype-matched controls (both R&D Systems, 20µg/ml unless otherwise stated). DC were nucleoporated with 5µg lentiviral plasmid DNA using the Amaxa Mouse Dendritic Cell Nucleofector Kit (Lonza). Unsorted cells were used for experiments. Splenic CD4⁺ T cells were isolated using a negative selection magnetic bead isolation kit (Miltenyi Biotech). For proliferation experiments, CD4⁺ T cells were labeled with 5µM CFSE dye for 20 min at 37°C then washed before co-culture.

ELISA and flow cytometry assays

Supernatants of LPS-stimulated DC were assayed for IL-1 β , IL-10, IL-12 using the Beadlyte[®] system (Millipore) and for IL-6 and TNF- α on ELISA (eBioscience). IL-2 secretion by CD4⁺ T cells was analysed with mouse IL-2 ELISA kit (R&D Systems). pSTAT5 assays were performed as previously described²⁵ using serum-starved CD4⁺ T cells co-cultured at 1:1 ratio for 10 min at 37°C with DC. Antibodies used for flow cytometry were against CD16/CD33 (2.4G2), CD86 (GL1), CD11c (HL3), CD4 (RM 4-5, SK3), I-A/I-E (2G9), pSTAT5 (pTyr694; clone 47) (all BD Biosciences), IL-15R α (AF551) (R&D systems) and against γ c (M-20) and IL-15 (H-114) (both from Santa Cruz Biotechnology). Apoptosis was assessed using the AnnexinV Apoptosis Detection Kit (BD Biosciences).

Antigen uptake and presentation assays

Uptake and breakdown of DQ-OVA (self-quenched fluorescent conjugate of ovalbumin, Molecular Probes, Invitrogen), measured as emission of green fluorescence (515nm), were assessed as previously described²⁴. For measurement of antigen presentation, DC were matured overnight with LPS in the absence or presence of the indicated concentrations of E α -GFP protein (kindly provided by Dr. Paul Garside, University of Glasgow). E α peptide presentation was measured after 24hrs by flow cytometry. Briefly, cells were stained with antibodies against CD11c, IA/IE and the biotinylated Yae (specific for E α ⁵²⁻⁶⁸ peptide presented on I-Ab) antibody (eBioscience) followed by streptavidin. DC were gated as CD11c⁺IA/IE^{hi} cells and presentation of E α calculated as an index relative to DC matured in the absence of E α -GFP (LPS only) using the following equation: $100 \times (\log^{E\alpha}/\log^{LPS}) - \log^{LPS}$. For measurement of antigen presentation, DC were pulsed overnight with varying concentrations of OVA in the presence of LPS then co-cultured for 48hrs at a 1:5 ratio with BO17.4 hybridoma cells. IL-2 secretion by BO17.4 cells was measured by ELISA.

Planar lipid bilayers

Liposome stocks containing DOPC, 25 mol% DGS-NTA and 2 mol% Cap-biotin (Avanti Polar Lipids) were prepared as described elsewhere²⁶. To make glass-supported planar bilayers for DC imaging, liposomes were mixed in appropriate ratios to produce DOPC bilayers with 0.01 mol% Cap-biotin and 12.5% DGS-NTA. Following washing with HBS containing 1% human serum albumin, 1mM Ca and 2mM Mg (HBS/HSA), bilayers were blocked with 5% Casein containing 100 μ M NiCl₂, and incubated with 5 μ g/ml streptavidin in HBS/HSA for 15min, and following washing, incubated for a further 30 min with a mixture of LFA-I domain-His6 (10 μ g/ml) and monobiotinylated anti-I-A/E Fab' fragments (5 μ g/ml). Further details of anti-I-A/E, LFA and ICAM protein preparations are available in the [Supplemental material](#). For imaging of OTII T cells, DOPC bilayers were prepared as above, containing 12.5% DOGS-NTA. ICAM-his12 and I-A^b/OVA-his12 were added to bilayers to yield densities of 300 mol/ μ m² and 100 mol/ μ m² respectively. DOPC liposomes containing CD80 were incorporated at 200 mol/ μ m². Soluble IL-15/IL-15R α with a C-terminal 6-histidine tag (eBioscience) was incorporated (2 μ g/ml) as indicated.

Microscopy

TIRF imaging was performed using a Nikon Ti microscope equipped with a 100x Nikon TIRF objective, NA 1.49. Cells interacting with bilayers were fixed with 2% PFA; permeabilised with 0.1% saponin and quenched with 50mM glycine; blocked and stained with pSTAT5 (D47E7) (Cell Signaling) or IL-15R α (H-107) (Santa Cruz Biotechnology). Secondary antibodies used were anti-rabbit AlexaFluor488 (Molecular Probes, Invitrogen). Measurement of labeled molecules was achieved by determining fluorescence intensity within regions of cell contact identified either using a threshold on TCR intensity (pSTAT5) or by the IRM channel (IL-15R α). For analysis of MHCII and GFP accumulation at DC interfaces, fluorescence intensities were acquired at 4 frames/min over 25-min. Data were analysed with the Metamorph and ImageJ software. Please see [Supplementary File 1](#) for more details on imaging methodology.

Imaging of DC on lipid bilayers

Tracking of DC by confocal imaging was performed at 37°C in a heated environmental chamber. LPS/OVA-stimulated DC were introduced into flow-cells and areas of bilayers, selected at random, imaged for 37–45 min at 15 sec intervals. DIC and reflection (IRM) channels were recorded (+/- AF568 fluorescence) using appropriate laser excitation and emission filters. Cells were tracked manually in ImageJ software using cell nuclei in DIC images as a position reference. For quantitation of fluorescence intensities at DC interfaces with planar bilayers by TIRFM, cell contacts in the central region of the TIRF field, which is more evenly illuminated than the edges, were analyzed to minimize variations due to the inherent curvature of TIRF mode illumination. To estimate the extent to which variations in TIRF illumination contributed to the observed differences in measurements of specific fluorescence, the anti-MHC II Fab' AF568 fluorescence intensity in bilayer regions immediately adjacent to DC interfaces was measured for all interfaces from which IL-15R α fluorescence intensity was quantitated. Since non-interface anti-MHC II Fab' AF568 is evenly distributed on bilayers, its fluorescence effectively represents laser excitation, in TIRF mode, within the imaging field. The morphology of the TIRF field was comparable between fluorescence channels. This baseline anti-MHC II Fab' AF568 fluorescence was used to estimate the contribution of inter-sample (between γ c^{-/-} and WT DC samples) variation in TIRF illumination in interface fluorescence intensity measurements. Colocalization between engaged MHC II and IL-15R α at DC interfaces was measured using Pearson correlation coefficient (PCC). To rule out spurious differences in PCC due to lower IL-15R α fluorescence intensity at γ c^{-/-} DC interfaces, PCC between MHC II and IL-15R α was calculated for a subset of γ c^{-/-} and WT interfaces with comparable IL-15R α fluorescence intensity.

Intracellular Ca²⁺ imaging

Bilayers containing LFA-1 I α with or without anti-MHCII Fab' fragments were made in FCS II flow cells as described above. Prior to introduction of DCs, flow cells were equilibrated to 37°C in the heated environmental of an LSM510 confocal microscope. DCs were loaded with 3 μ M Fluo-4 AM (Invitrogen)

for 20 min in serum free media, washed, and incubated for a further 20 min in complete cell culture media. Cells were subsequently washed, resuspended in HBS/HSA and introduced into flow chambers for confocal imaging using a 20x, NA 0.75 air objective, and wide confocal iris settings. All imaging was performed at 37°C, and images acquired for Fluo-4 and DIC channels every 15 seconds for ~25 minutes. Cell tracking and mean Fluo-4 fluorescence was measured using ImageJ.

Statistics

Prism v.5 (GraphPad Software) was used for statistical analysis. This included two-tailed Student's t-test with 95% confidence bounds, one-way ANOVA (with Bonferroni's correction for multiple comparison), Gaussian curve-fitting was performed with single, bimodal, and trimodal model parameters.

Results

$\gamma\text{c}^{-/-}$ DC fail to trans-present IL-15 during antigen specific activation of naïve CD4+ T cells

To investigate the role of γc in DC function, we generated conventional bone marrow-derived DC (BMDC) from γc -deficient ($\gamma\text{c}^{-/-}$) mice²⁷. These mice also lack lymphoid-restricted recombinase activating gene 2 (RAG 2) by genetic modification, to eliminate low levels of persisting T cells seen in γc single knockout strains²⁸. Deletion of RAG 2 does not impair the function of GM-CSF-derived BMDC²⁹ consistent with a lack of expression of VDJ rearrangement genes and RAG transcripts in conventional DC³⁰.

As expected, BMDC derived from $\gamma\text{c}^{-/-}$ mice completely lacked γc protein expression and γc -dependent cytokine signaling (Figure S1A,B) but expressed MHCII and the costimulatory molecule CD86 at levels comparable with WT DC both in the immature state and following LPS-induced maturation (Figure S1C). To test whether $\gamma\text{c}^{-/-}$ DC support normal antigen-mediated priming of T cells, DC were pulsed with whole ovalbumin (OVA) that is internalized, processed and presented on the surface of DC as a peptide antigen (pOVA) in complex with the class II MHC molecule I-A^b. When co-cultured with OTII CD4+ T cells, transgenic for a TCR recognising pOVA/I-A^b, OVA-pulsed $\gamma\text{c}^{-/-}$ DC induced a moderate but significantly lower level of T cell proliferation than WT DC ($p \leq 0.05$, Figure 1A,B) and markedly reduced IL-2 secretion ($p \leq 0.05$, Figure 1C). As previously described³¹, under these conditions, IL-2 release by DC was negligible (Figure S1D) indicating that the impairment was due to defective T cell activation. Taken together, these data show that DC-expressed γc is required for full activation of antigen-specific CD4+ T cells.

Our findings were not due to impaired antigen uptake, processing or presentation of surface MHC/antigen complexes as $\gamma\text{c}^{-/-}$ DC were as efficient as WT DC at internalising and processing DQTM ovalbumin (Figure S1E) and at processing and presenting the model E α antigen (Figure S1F,G). Furthermore, OVA-pulsed mature $\gamma\text{c}^{-/-}$ and WT DC induced similar levels of IL-2 release from the B017.4 T cell hybridoma (which expresses the OTII TCR and is less dependent on costimulation) (Figure S1H), demonstrating that pOVA presentation by surface MHC molecules was

functionally similar between WT and $\gamma\text{c}^{-/-}$ DC. Together, these data demonstrate that the observed defects in $\gamma\text{c}^{-/-}$ DC mediated CD4+ T cell activation are not explained by defective antigen handling. As mature $\gamma\text{c}^{-/-}$ and WT DC released similar levels of pro-inflammatory cytokines such as IFN- β , IL6, IL-12 and TNF- α (Figure S1I), we reasoned that the observed defect of T-cell activation was due to a contact dependent rather than a soluble messenger mechanism.

As it is known that optimal naïve T cell activation depends on stable adhesion to DC³², we investigated whether γc -deficiency impaired T-DC intercellular adhesion. Both conjugate-formation and redistribution of LFA-1 to the IS, a hallmark of T cell polarisation in response to antigen recognition³³, were preserved in T cells co-cultured with $\gamma\text{c}^{-/-}$ DC (Figure S2A–C). Taken together, these data demonstrate that the defective antigen-specific T cell priming observed in $\gamma\text{c}^{-/-}$ DC is not due to impaired adhesion or LFA-1/ICAM-1 dependent T cell polarization. We further examined the fine-structure of the T-DC contact interface using transmission electron microscopy. Binding of TCR to pMHC occurs at, and stabilises, regions of close contact (~12 nm apart) between apposed membranes at T-DC interfaces, which are thought to be critical for signaling^{34,35}. Compared to WT DC interfaces, $\gamma\text{c}^{-/-}$ DC formed a similar proportion of close contacts with T cells, interspersed between areas of greater membrane separation (~30–50 nm)(Figure S2D–G), demonstrating that antigen-induced close contacts were preserved in the absence of γc .

Since antigen presentation, adhesion, secretory, and canonical costimulatory functions appeared to be preserved in $\gamma\text{c}^{-/-}$ DC, we considered other plausible defects in DC function that might account for incomplete T cell priming. One candidate for this is the delivery of IL-15 mediated stimulatory signals to T cells by transpresentation. While it is well established that DC trans-present IL-15 to CD8+ T cells and NK cells^{36,37}, the role of IL-15 transpresentation in CD4+ T cell activation has only begun to be explored²³. To further establish whether IL-15 transpresentation occurs during DC-CD4+ T cell interactions, we analysed STAT5 activation in OTII T cells after 10 minute co-culture with LPS-matured WT DC. We observed that significant induction of STAT5 phosphorylation in CD4+ T cells occurred only when DC had been pre-loaded with antigen ($p \leq 0.05$, Figure 1D,E). Pre-treatment with an IL-15 blocking (Figure 1D,E), but not an IL-2 blocking (Figure S3A,B), monoclonal antibody abolished STAT5 phosphorylation in OTII T cells indicating that STAT5 activation occurs primarily through IL-15 transpresentation during DC-mediated priming of naïve CD4+ T cells in our experimental system.

Notably, antigen-pulsed $\gamma\text{c}^{-/-}$ DC were severely compromised in their ability to activate STAT5 in OTII T cells, compared with WT DC (Figure 1D,E), strongly implicating a role for γc in IL-15 transpresentation by DC. T cell proliferation induced by antigen-pulsed WT DC was also inhibited by IL-15R α blockade in a dose-dependent manner. Consistent with the notion that naïve CD4+ T cell priming by $\gamma\text{c}^{-/-}$ DC is compromised primarily due to defective IL-15 transpresentation, the reduced antigen-specific




Figure 1. $\gamma^{\text{c-/-}}$ DC fail to transpresent IL-15 to CD4⁺ T cells during antigen-specific priming. (A) Representative plots showing CFSE dilution in CD4⁺-gated OTII T cells after co-culture with LPS-matured OVA-pulsed DC for 72 hrs. Grey dotted histograms represent CD4⁺ T cells cultured with DC in the absence of antigen. (B) Quantification of T cell proliferation shown in A. (C) IL-2 release by CD4⁺ T cells incubated with OVA-pulsed DC at the indicated ratios for 72hrs. (D) pSTAT5 induction in CD4⁺ T cells following a 10 minute incubation with DC, either untreated or stimulated overnight with LPS $\pm$ OVA. DC were blocked with anti-IL-15R α or isotype control. (E) Increase in pSTAT5 fluorescence (D), compared to unstimulated control. (F) Inhibition of CD4⁺ T cell proliferation after co-culture for 72hr with OVA-pulsed DC pre-treated with anti-IL-15R α (relative to isotype-matched control antibody treatment). (G) Total and surface IL-15R α expression (FAB551F) on CD11c gated cells. (H) Schematic of construct encoding full length (γ^{WT} -GFP) and truncated ($\gamma^{\Delta\text{C}}$ -GFP) γ^{c} attached to GFP. (I) Increase in pSTAT5 levels, compared to unstimulated control, in CD4⁺ T cells following incubation with DC $\pm$ OVA, transfected with γ^{WT} -GFP, $\gamma^{\Delta\text{C}}$ -GFP or ctrl GFP. *P* values, *t*-test (B,C,I); one-way ANOVA (E); linear regression (*p* value tests for significant difference between the slope of each line) (F).

proliferative response of naïve CD4⁺ T cells to $\gamma_c^{-/-}$ DC was not further affected by IL-15R α blockade ($p \leq 0.01$, [Figure 1F](#) and [Figure S3C](#)). The observed differences in T cell proliferation were not attributable to differential post-activation T cell viability ([Figure S3D](#)), or IL-15R α expression, as both total and surface levels of IL-15R α were comparable between WT and $\gamma_c^{-/-}$ DC ([Figure 1G](#)). Levels of total and surface IL-15 available for transpresentation were also unaffected by absence of DC- γ_c ([Figure S3E](#)).

To more definitely establish whether γ_c expression in DC was necessary for IL-15 trans-signaling to T cells, we transfected $\gamma_c^{-/-}$ DC with constructs encoding either GFP alone, the full-length γ_c fused to GFP³⁸ (γ_c^{WT} -GFP), or a truncated γ_c ($\gamma_c^{\Delta c}$ -GFP), that lacks 77 amino acids at the cytoplasmic carboxy-terminus, allowing surface expression but not signaling function³⁹ ([Figure 1H](#) and [Figure S3F,G](#)). Expression of γ_c^{WT} -GFP or $\gamma_c^{\Delta c}$ -GFP in $\gamma_c^{-/-}$ DC rescued STAT5 activation in OTII T cells, while expression of GFP alone had no effect ([Figure 1I](#) and [Figure S3H](#)). Similar levels of GFP expression ($\approx 35\%$) were obtained with all constructs ([Figure S3I](#)).

To confirm that IL-15 trans-presentation at the CD4⁺ T cell IS requires MHC:TCR engagement, we employed glass-supported planar lipid bilayers containing ICAM-1, CD80 and pOVA/I-A^b to recapitulate the essential features of an antigen-presenting surface suitable for naïve T cell stimulation, and incorporated

IL-15/IL-15R α complexes to mimic DC-mediated IL-15 transpresentation. Using this model system, we measured STAT5 phosphorylation as a marker of IL-2R β/γ_c mediated trans-signaling at the T cell IS by TIRFM. As expected, naïve OTII T cells formed a mature IS, at which TCR accumulated in a central supramolecular cluster (cSMAC)³⁹, only in response to pOVA/I-A^b ([Figure 2A](#), arrows). In keeping with our flow-cytometry results, IL-15/IL-15R α did not activate STAT5 signaling in the absence of antigen, suggesting that TCR engagement is required for IL-15/IL-15R α mediated trans-signaling in CD4⁺ T cells ([Figure 2A,B](#) and [Figure S4A,B](#)). Despite dependence on TCR engagement for IL-15/IL-15R α mediated STAT5 activation, PLC γ 1 phosphorylation, which occurs downstream of TCR/CD28 signaling, and Akt phosphorylation, which is strongly induced by CD28 ligation, were not affected by IL-15 trans-signaling ([Figure S5A–D](#)). Similarly, Zap-70 phosphorylation following TCR triggering was not affected by IL-15 trans-signaling, indicating minimal cross-talk between TCR/CD28 signals and the JAK/STAT pathway ([Figure S5E–G](#)).

Taken together, these data demonstrate that DC transpresent IL-15 to CD4⁺ T cells by a mechanism that depends on MHCII/TCR ligation and DC-expressed γ_c . Surprisingly, γ_c signaling in DC was not required for IL-15 transpresentation, suggesting that γ_c facilitates IL-15 transpresentation through induced intermolecular *cis* interactions and/or cytoskeletal reorganization at the intramembrane or ectodomain level.




Figure 2. IL-15 mediated trans-signaling in CD4⁺ T cells requires TCR engagement. (A) Representative TIRFM images of CD4⁺ T cells incubated on glass-supported planar bilayers containing ICAM-1, CD80, OVA₃₂₃₋₃₃₉/I-A^b and IL-15/IL-15R α , as indicated. After 30 min incubation at 37°C in HBS/HSA buffer, cells were fixed and stained for TCR (red, Alexa Fluor 568) and phospho-STAT5 (pSTAT5) (green, Alexa Fluor 488) in PBS buffer. White arrows in the image panels indicate central accumulation of TCR at the T cells IS. (B) Quantification of pSTAT5 fluorescence in A. Data are presented as percentage increase in pSTAT5 fluorescence, relative to unstimulated controls. Data are from 2 independent experiments (N=32-57) (mean $\pm$ S.E.M). *P* values, one-way ANOVA. Imaging was performed on a Nikon Ti microscope with a 100x TIRF objective, N.A. 1.49, controlled by Nikon Elements software. Fluorescence images were captured using an Ixon cooled EMCCD camera (512 x 512 pixels, Andor Technology). Mean fluorescence intensity at contact interfaces was quantified from 14 bit images using Metamorph software. Brightness and contrast are adjusted uniformly across image groups for clarity.

Binding-induced clustering and accumulation of MHCII at the DC IS

To investigate molecular events that follow MHC ligation at the DC IS at high spatial resolution, we developed a glass-supported bilayer system that recapitulates both adhesive and MHC-binding properties of the T cell surface (Figure 3A). To ligate ICAM-1 on DC, we loaded bilayers with a C-terminally 6 histidine tagged inserted domain fragment of the LFA-1 α -subunit (α I) that is covalently locked in its high affinity conformation⁴⁰. To ligate MHCII, we generated C-terminally monobiotinylated Fab' fragments⁴¹ of an I-A/E specific monoclonal antibody (clone M5/114) to approximate TCR ectodomain size and valency (see Supplementary methods). These surrogate TCRs were attached to bilayers containing biotin headgroups *via* a streptavidin 'bridge', ensuring a uniform orientation that is favourable for MHC binding. Fragments were labelled with fluorophores (f/p $\sim$ 3) to follow their recruitment and lateral reorganization upon binding to MHCII at the DC contact interface by confocal and TIRF microscopy.

Initial imaging by confocal microscopy revealed that WT DC exhibit a 'crawling' motility (mean velocity $\sim$ 8 μ m/min) on bilayers containing LFA-1 (Movie S1, Figure S6A,C). Ligation of MHCII on WT DC led to an arrest in motility (mean velocity $\sim$ 2.5 μ m/min) and accumulation of engaged MHCII at the DC-bilayer interface (Movie S2, Figure S6A,C). Although γ c^{-/-} DC migrated more slowly (mean velocity $\sim$ 6 μ m/min) compared to WT DC, ligation of MHCII led to a similar arrest in motility (Movie S3, Figure S6B,D), indicating that MHCII ligation delivers a 'stop' signal to DCs, analogous to that in T cells following antigen recognition⁴², that is not dependent on DC- γ c expression. MHC II ligation does not lead to a rise in intracellular Ca²⁺ levels in WT and γ c^{-/-} DC (Movie S4,5 and Figure S6E-G), suggesting that, in contrast to antigen-induced T cell stopping, motility arrest following MHC II ligation in DC is not associated with Ca²⁺ signaling.

We next investigated the binding-induced organization of MHC II at the DC IS by total internal reflection fluorescence microscopy (TIRFM). Within seconds of contact with bilayers, ligated MHCII formed small clusters throughout the contact interface (Figure 3B and Figure S6H), which were transported towards the center of the contact interface, presumably by interaction with the DC cytoskeleton²⁴ (Movie S6,7). The extent of MHC accumulation at the contact interface was similar to that of WT DC (Figure S6H,I).

DC-expressed γ c controls IL-15R α recruitment to the IS

Mirroring MHC polarisation to the IS^{43,44}, γ c^{WT}-GFP and γ c^{Δc}-GFP expressed in γ c^{-/-} DC were recruited to the contact interface, leading to $\sim$ 4-fold enrichment of mean GFP fluorescence over 23 minutes (Figure 3B,E and 3C,F). In contrast, GFP alone was not enriched at the contact interface over the same time course (Figure 3D,G).

Since DC-expressed γ c was critical for effective IL-15 transpresentation in co-culture assays, its recruitment and colocalization with MHCII at the DC IS suggested the possibility of a

spatially regulated mechanism for transpresentation, in which γ c coordinates recruitment of IL-15/IL-15R α to the DC synapse following MHCII ligation. To test this hypothesis, we imaged WT DC on bilayers at an early time-point (15 minutes), in the presence or absence of surrogate TCRs (anti-I-A/E Fab'), and labelled IL-15R α for TIRF imaging. When compared to bilayers containing only LFA-1 α I domain, ligation of MHCII induced almost 3-fold more IL-15R α at the DC IS, demonstrating that MHCII engagement is sufficient to recruit IL-15R α to the IS in WT DC (Figure 4A,B). Strikingly, MHCII-induced IL-15R α recruitment to the IS was severely compromised in γ c^{-/-} DC (p $\leq$ 0.001, Figure 4A,B) while MHCII accumulation was relatively unaffected (Figure 4C). These differences were not accounted for by variations in TIRF imaging conditions between samples (Figure S7). The extent of colocalization between MHCII and IL-15R α was also decreased in the absence of γ c, suggesting that it promoted a closer association between engaged MHCII and IL-15R α (Figure 4D and Figure S8). Expression of γ c^{WT}-GFP and γ c^{Δc}-GFP in γ c^{-/-} DC also resulted in $\sim$ 70% increase in IL-15R α accumulation at the interface upon MHC II ligation, when compared to expression of GFP alone (Figure 5A,B), indicating that γ c signaling in DC was dispensable for IL-15R α recruitment.

Discussion

Patients with γ c-deficient SCID remain susceptible to opportunistic HPV infections even when T cell function is restored by BMT. This raises the possibility that residual γ c-deficient DC, which persist in the absence of myeloablative conditioning, might be ineffective in priming T cell immunity. To investigate this in a tractable model, we tested the ability of bone-marrow derived DC from γ c knockout mice to activate normal naïve CD4+ T cells. We have identified defects in the ability of γ c-deficient DC to activate antigen-specific CD4+ T cells, which could not be accounted for by a problem with DC maturation or antigen processing. Instead, our studies have revealed an unexpected requirement for IL-15 transpresentation in CD4+ T cell activation. Furthermore, we have identified a role for DC γ c in the recruitment of IL-15R α to the DC side of the immune synapse, which is critical for effective IL-15 transpresentation to CD4+ T cells, and is independent of γ c signaling function. Therefore, our *in vitro* functional and imaging studies have revealed a mechanism that may account for a subset of immune dysfunction in γ c-deficient myeloid cells. While these studies have generated new hypotheses that can be explored further in human DC, the finding that IL-15 transpresentation contributes to CD4+ T cell activation in a DC γ c-dependent manner, extends our understanding of the costimulatory requirements for CD4+ T cell priming. High resolution imaging of DC using a planar bilayer model system has provided new perspectives on the active role of DC in IS formation, that we expect will be useful for further investigation of the DC IS.

Soluble IL-15 produced in DC binds effectively irreversibly (K_d $\sim$ 10⁻¹¹ M)⁴⁵ to co-expressed IL-15R α within intracellular compartments, before trafficking to the DC cell surface, for transpresentation to T and NK cells expressing IL-2R β / γ c heterodimers^{46,47}. Signal transduction in T cells occurs through the cytoplasmic portions of IL-2R β / γ c heterodimers, which are




Figure 3. γ is recruited with MHCII at the DC IS. (A) Schematic diagram of a glass-supported planar bilayer recapitulating a T cell surface. (B–D) Representative TIRFM images demonstrating GFP (green) and MHCII (red, Alexa Fluor 568) accumulation over time at the contact interface of LPS/OVA stimulated $\gamma^{-/-}$ DC expressing γ^{WT} -GFP (B), $\gamma^{\Delta C}$ -GFP (C) or GFP alone (D), interacting on bilayers shown in A. (E–G) Time course of GFP accumulation at the contact interface in DC transfected with γ^{WT} -GFP (E) (N=6), $\gamma^{\Delta C}$ -GFP (F) (N=5) or GFP alone (G) (N=5). Data represent mean fluorescence intensity (mean+S.E.M. are shown for clarity), normalized relative to values at initial point of contact by DC on bilayers (t=0). Live cells in HBS/HSA buffer were imaged in FCS2 flow chambers (Biopetech) maintained at 37°C. Imaging was performed on a Nikon Ti microscope with a 100x TIRF objective, N.A. 1.49, controlled by Nikon Elements software. Fluorescence images were captured using an Ixon cooled EMCCD camera (512 x 512 pixels, Andor Technology). Mean fluorescence intensity at contact interfaces was quantified from 14 bit images using Metamorph software. Brightness and contrast are adjusted uniformly across image groups for clarity.




Figure 4. IL-15R α is recruited to the DC interface and colocalizes with engaged MHCII. LPS/OVA stimulated DC were incubated on glass-supported planar bilayers containing LFA-1 α I ± I-A/E Fab'. After 15 min incubation at 37°C in HBS/HSA buffer, cells were fixed, permeabilized and stained for IL-15R α (Alexa Fluor 488) in PBS buffer. **(A)** Representative TIRFM images showing IL-15R α and MHCII (Alexa Fluor 568) accumulation at contact interfaces. Fluorescence intensity profiles (right panels) indicate the distribution of IL-15R α (green) and MHCII (red) along the dashed white lines (merge). **(B,C)** Quantification in arbitrary units (AU) of mean IL-15R α fluorescence **(B)** and MHCII fluorescence **(C)** at DC contact interfaces shown in A (N=89-90, mean $\pm$ S.E.M). **(D)** Quantification of colocalization between MHCII and IL-15R α at DC contact interfaces **(A)**, calculated as Pearson's correlation coefficient (PCC). Imaging was performed on a Nikon Ti microscope with a 100x TIRF objective, N.A. 1.49, controlled by Nikon Elements software. Fluorescence images were captured using an Ixon cooled EMCCD camera (512 x 512 pixels, Andor Technology). Mean fluorescence intensity at contact interfaces was quantified from 14 bit images using Metamorph software. PCC was calculated for MHCII and IL-15R α fluorescence channels using ImageJ software. Brightness and contrast are adjusted uniformly across image groups for clarity.

associated with Janus family tyrosine kinases JAK1 and JAK3. Assembly of the IL-15/IL-15R ternary complex in *trans* leads to activation of JAK1/3, and subsequent phosphorylation of IL-2R β / γ_C . This leads to recruitment and activation of signal

transducer and activator of transcription 5 (STAT5) proteins¹². We have shown in functional studies that IL-15 transpresentation by DC to CD4⁺ T cells is critically dependent on DC-expressed γ_C . Our imaging studies demonstrate that MHCII ligation leads to




Figure 5. γ_C is required for IL-15R α recruitment to the DC IS. (A) Representative TIRFM images showing GFP, MHCII (Alexa Fluor 568) and IL-15R α (Alexa Fluor 633) accumulation at contact interfaces of $\gamma_C^{-/-}$ DC, transfected with γ_C^{WT} -GFP, $\gamma_C^{\Delta C}$ -GFP or ctrl GFP. **(B)** Quantification in arbitrary units (AU) of mean IL-15R α fluorescence at contact interfaces shown in A (N=38-43, mean $\pm$ S.E.M). P values, one-way ANOVA. After incubation on bilayers at 37°C in HBS/HSA buffer, cells were fixed and permeabilized to stain for IL-15R α in PBS buffer. Imaging was performed on a Nikon Ti microscope with a 100x TIRF objective, N.A. 1.49, controlled by Nikon Elements software. Fluorescence images were captured using an Ixon cooled EMCCD camera (512 x 512 pixels, Andor Technology). Mean fluorescence intensity at contact interfaces was quantified from 14 bit images using Metamorph software. Brightness and contrast are adjusted uniformly across image groups for clarity.

γ_C dependent recruitment of IL-15R α to the DC IS, where it colocalizes with engaged MHCII. Both IL-15R α recruitment to the DC IS, and IL-15 mediated trans-signaling in CD4⁺ T cells, are restored in γ_C -deficient DC following re-expression of γ_C . Neither process appeared to depend on signaling function as truncation of the γ_C cytoplasmic tail was also effective in recruiting IL-15R α . Curiously, transpresented IL-15 triggered STAT5 signaling in CD4⁺ T cells only when TCR was engaged.

A precise picture of the molecular dynamics and subunit stoichiometries, in cell membranes, of IL-15/IL-15R α and its associated receptor subunits has not yet been established. However, elegant imaging studies of IL-15R α in transformed and primary T cell lines have revealed considerable heterogeneity in subunit composition, and a far more diverse set of *cis* associations, than might be predicted by 'affinity conversion' or other assembly models⁴⁸. Of relevance to our findings, MHCII has been shown to associate with both IL-15R α ⁴⁹ and with γ_C ⁵⁰. Our observations, that γ_C is recruited to the DC IS, plays a critical role in recruitment of IL-15R α , and promotes greater colocalization between engaged MHCII and IL-15R α , lead us to favor a molecular configuration on the DC cell surface in which IL-15R α , γ_C , and MHCII exist as a loosely coupled molecular complex, that is consolidated by MHCII engagement. Since MHCII engagement leads to its clustering and dynamic

transport, presumably by interaction with the DC cytoskeleton, MHCII-nucleated domains may serve as avidity-enhancing scaffolds, or platforms within liquid-ordered lipid domains⁵¹ that stabilize the IL-15R α - γ_C -MHCII trimolecular association.

Taken together, our findings suggest a model of IL-15 transpresentation in which peptide/MHCII ligation by cognate TCR results in γ_C -mediated recruitment of IL-15R α (in complex with IL-15) to the DC synapse, where it is positioned near sites of TCR engagement for binding *in trans* (Figure 6A,B). Coupled delivery of IL-15 mediated costimulation with antigen recognition is consistent with the suggestion that close membrane apposition at the DC-T cell interface, determined by the (small) size of pMHC/TCR and accessory receptor complexes (~15 nm), may favor assembly of the *trans* IL-15/IL-15R ternary complex at DC-T-cell interfaces, since it is similar in size to TCR/pMHC complexes⁵². TCR ligation of pMHC drives the formation of close contacts with APC, from which the large T cell surface phosphatase CD45, a key negative regulator of both TCR⁵³, and γ_C cytokine receptor-associated JAK signaling⁵⁴, is excluded^{35,55}. Coupled (trans)presentation of pMHC and IL-15 at the DC-T cell IS may therefore allow spatially coordinated activation of the biochemically distinct TCR and JAK/STAT signaling pathways during antigen-specific priming of naïve T cells by DC.




Figure 6. Model of γ c-facilitated IL-15/IL-15R α transpresentation to CD4⁺ T cells. (A) In the absence of cognate peptide-MHC II (pMHCII), IL-2R β / γ c cytokine receptor signaling in T cells is not initiated (open circles denote unphosphorylated IL-2R β / γ c and associated JAK1/3), as these small receptors are likely positioned too far apart for stable binding of IL-15/IL-15R α complexes on the DC surface. Inset depicts putative association of IL-15/IL-15R α , MHCII and γ c within liquid-ordered (Lo) lipid domains, and/or through cytoskeletal confinement. (B) Engagement of pMHCII on DC (1) leads to γ c-dependent recruitment IL-15/IL-15R α to the contact interface, close to regions of bound pMHCII (2); this would position IL-15/IL-15R α complexes on the DC surface at a distance compatible with binding IL-2R β / γ c receptors in *trans* (2). Close contacts also exclude CD45, allowing stable phosphorylation (red circles) of both TCR and IL-2R β / γ c receptors, permitting recruitment and phosphorylation of STAT5 (3) in the context of productive antigen recognition.

Data availability

Dataset 1: *Dendritic cell-expressed common gamma-chain recruits IL-15 for trans-presentation at the murine immunological synapse* is available from OSF: <https://doi.org/10.17605/OSF.IO/YC7WS>⁵⁶.

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

Please see [Supplementary File 4](#) for the data legend. Image data are available on request, see person to contact in [Supplementary File 4](#).

Competing interests

No competing interests were disclosed.

Grant information

This work was supported by Wellcome Trust [090233, GB, AJT] and [100262Z, MLD]; the Child Health Research Appeal Trust and a Bogue Research Fellowship (CB); the Primary Immunodeficiency Association and the ICH Biomedical Research Centre (SOB); The European Union (GB); The Cancer Research Institute (KC); and US National Institutes of Health [grant R37 AI43542, MLD] and [R01 CA31798, TAS].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Supplementary materials

Supplementary File 1: Supplementary methods.

[Click here to access the data.](#)

Supplementary File 2: Supplementary figures (Figure S1–8).

[Click here to access the data.](#)

Supplementary File 3: Supplementary movies (Movie S1–7).

[Click here to access the data.](#)

Supplementary File 4: Data legend.

[Click here to access the data.](#)

References

- Gaspar HB, Harwood C, Leigh I, *et al.*: **Severe cutaneous papillomavirus disease after haematopoietic stem-cell transplantation in patients with severe combined immunodeficiency.** *Br J Haematol.* 2004; **127**(2): 232–233.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Laffort C, Le Deist F, Favre M, *et al.*: **Severe cutaneous papillomavirus disease after haematopoietic stem-cell transplantation in patients with severe combined immune deficiency caused by common gamma cytokine receptor subunit or JAK-3 deficiency.** *Lancet.* 2004; **363**(9426): 2051–2054.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Abd Hamid IJ, Slatter MA, McKendrick F, *et al.*: **Long-term outcome of hematopoietic stem cell transplantation for IL2RG/JAK3 SCID: a cohort report.** *Blood.* 2017; **129**(15): 2198–2201.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Belz GT, Nutt SL: **Transcriptional programming of the dendritic cell network.** *Nat Rev Immunol.* 2012; **12**(2): 101–113.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fahey LM, Raff AB, Da Silva DM, *et al.*: **Reversal of human papillomavirus-specific T cell immune suppression through TLR agonist treatment of Langerhans cells exposed to human papillomavirus type 16.** *J Immunol.* 2009; **182**(5): 2919–2928.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nakayama Y, Asagoe K, Yamauchi A, *et al.*: **Dendritic cell subsets and immunological milieu in inflammatory human papilloma virus-related skin lesions.** *J Dermatol Sci.* 2011; **63**(3): 173–183.
[PubMed Abstract](#) | [Publisher Full Text](#)
- de Jong A, van der Burg SH, Kwappenberg KM, *et al.*: **Frequent detection of human papillomavirus 16 E2-specific T-helper immunity in healthy subjects.** *Cancer Res.* 2002; **62**(2): 472–479.
[PubMed Abstract](#)
- Steele JC, Mann CH, Rookes S, *et al.*: **T-cell responses to human papillomavirus type 16 among women with different grades of cervical neoplasia.** *Br J Cancer.* 2005; **93**(2): 248–259.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Stanley M: **Immune responses to human papillomavirus.** *Vaccine.* 2006; **24** Suppl 1: S16–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Dillon S, Sasagawa T, Crawford A, *et al.*: **Resolution of cervical dysplasia is associated with T-cell proliferative responses to human papillomavirus type 16 E2.** *J Gen Virol.* 2007; **88**(Pt 3): 803–813.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mbulawa ZZ, Marais DJ, Johnson LF, *et al.*: **Influence of human immunodeficiency virus and CD4 count on the prevalence of human papillomavirus in heterosexual couples.** *J Gen Virol.* 2010; **91**(Pt 12): 3023–3031.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rochman Y, Spolski R, Leonard WJ: **New insights into the regulation of T cells by gamma(c) family cytokines.** *Nat Rev Immunol.* 2009; **9**(7): 480–490.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Koka R, Burkett P, Chien M, *et al.*: **Cutting edge: murine dendritic cells require IL-15R alpha to prime NK cells.** *J Immunol.* 2004; **173**(6): 3594–3598.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wuest SC, Edwan JH, Martin JF, *et al.*: **A role for interleukin-2 trans-presentation in dendritic cell-mediated T cell activation in humans, as revealed by daclizumab therapy.** *Nat Med.* 2011; **17**(5): 604–609.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Olsen SK, Ota N, Kishishita S, *et al.*: **Crystal Structure of the interleukin-15.interleukin-15 receptor alpha complex: insights into trans and cis presentation.** *J Biol Chem.* 2007; **282**(51): 37191–37204.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Stonier SW, Ma LJ, Castillo EF, *et al.*: **Dendritic cells drive memory CD8 T-cell homeostasis via IL-15 transpresentation.** *Blood.* 2008; **112**(12): 4546–4554.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bulfone-Paus S, Dürkop H, Paus R, *et al.*: **Differential regulation of human T lymphoblast functions by IL-2 and IL-15.** *Cytokine.* 1997; **9**(7): 507–513.
[PubMed Abstract](#) | [Publisher Full Text](#)
- de Jong JL, Farner NL, Sondel PM: **Distinctions in lymphocyte responses to IL-2 and IL-15 reflect differential ligand binding interactions with the IL-2Rbeta chain and suggest differential roles for the IL-2Ralpha and IL-15Ralpha subunits.** *Cytokine.* 1998; **10**(12): 920–930.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Grabstein KH, Eisenman J, Shanebeck K, *et al.*: **Cloning of a T cell growth factor that interacts with the beta chain of the interleukin-2 receptor.** *Science.* 1994; **264**(5161): 965–968.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Korholz D, Banning U, Böning H, *et al.*: **The role of interleukin-10 (IL-10) in IL-15-mediated T-cell responses.** *Blood.* 1997; **90**(11): 4513–4521.
[PubMed Abstract](#)
- Dooms H, Desmedt M, Vancaeneghem S, *et al.*: **Quiescence-inducing and antiapoptotic activities of IL-15 enhance secondary CD4⁺ T cell responsiveness to antigen.** *J Immunol.* 1998; **161**(5): 2141–2150.
[PubMed Abstract](#)
- Purton JF, Tan JT, Rubinstein MP, *et al.*: **Antiviral CD4⁺ memory T cells are IL-15 dependent.** *J Exp Med.* 2007; **204**(4): 951–961.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Waickman AT, Ligons DL, Hwang S, *et al.*: **CD4 effector T cell differentiation is controlled by IL-15 that is expressed and presented in trans.** *Cytokine.* 2017; **99**: 266–274.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bouma G, Mendoza-Naranjo A, Blundell MP, *et al.*: **Cytoskeletal remodeling mediated by WASp in dendritic cells is necessary for normal immune synapse formation and T-cell priming.** *Blood.* 2011; **118**(9): 2492–2501.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zhang F, Thornhill SI, Howe SJ, *et al.*: **Lentiviral vectors containing an enhancer-less ubiquitously acting chromatin opening element (UCOE) provide highly reproducible and stable transgene expression in hematopoietic cells.** *Blood.* 2007; **110**(5): 1448–1457.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dustin ML, Starr T, Varma R, *et al.*: **Supported planar bilayers for study of the immunological synapse.** *Curr Protoc Immunol*, edited by John E Coligan [et al]. 2007; **Chapter 18**: Unit 18.13.
[PubMed Abstract](#) | [Publisher Full Text](#)
- DiSanto JP, Müller W, Guy-Grand D, *et al.*: **Lymphoid development in mice with a**

- targeted deletion of the interleukin 2 receptor gamma chain. *Proc Natl Acad Sci U S A*. 1995; **92**(2): 377–381.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Goldman JP, Blundell MP, Lopes L, *et al.*: Enhanced human cell engraftment in mice deficient in RAG2 and the common cytokine receptor gamma chain. *Br J Haematol*. 1998; **103**(2): 335–342.
[PubMed Abstract](#) | [Publisher Full Text](#)
 29. Faure M, Villiers CL, Marche PN: Normal differentiation and functions of mouse dendritic cells derived from RAG-deficient bone marrow progenitors. *Cell Immunol*. 2004; **228**(1): 8–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
 30. Shigematsu H, Reizis B, Iwasaki H, *et al.*: Plasmacytoid dendritic cells activate lymphoid-specific genetic programs irrespective of their cellular origin. *Immunity*. 2004; **21**(1): 43–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Granucci F, Feau S, Angeli V, *et al.*: Early IL-2 production by mouse dendritic cells is the result of microbial-induced priming. *J Immunol*. 2003; **170**(10): 5075–5081.
[PubMed Abstract](#) | [Publisher Full Text](#)
 32. Huppa JB, Gleimer M, Sumen C, *et al.*: Continuous T cell receptor signaling required for synapse maintenance and full effector potential. *Nat Immunol*. 2003; **4**(8): 749–755.
[PubMed Abstract](#) | [Publisher Full Text](#)
 33. Monks CR, Freiberg BA, Kupfer H, *et al.*: Three-dimensional segregation of supramolecular activation clusters in T cells. *Nature*. 1998; **395**(6697): 82–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
 34. Anton van der Merwe P, Davis SJ, Shaw AS, *et al.*: Cytoskeletal polarization and redistribution of cell-surface molecules during T cell antigen recognition. *Semin Immunol*. 2000; **12**(1): 5–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
 35. Choudhuri K, Wiseman D, Brown MH, *et al.*: T-cell receptor triggering is critically dependent on the dimensions of its peptide-MHC ligand. *Nature*. 2005; **436**(7050): 578–582.
[PubMed Abstract](#) | [Publisher Full Text](#)
 36. Lucas M, Schachterle W, Oberle K, *et al.*: Dendritic cells prime natural killer cells by trans-presenting interleukin 15. *Immunity*. 2007; **26**(4): 503–517.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 37. Stonier SW, Schluns KS: Trans-presentation: a novel mechanism regulating IL-15 delivery and responses. *Immunol Lett*. 2010; **127**(2): 85–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 38. Mussolino C, Morbitzer R, Lütge F, *et al.*: A novel TALE nuclease scaffold enables high genome editing activity in combination with low toxicity. *Nucleic Acids Res*. 2011; **39**(21): 9283–9293.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Morelon E, Dautry-Varsat A, Le Deist F, *et al.*: T-lymphocyte differentiation and proliferation in the absence of the cytoplasmic tail of the common cytokine receptor gamma c chain in a severe combined immune deficiency X1 patient. *Blood*. 1996; **88**(5): 1708–1717.
[PubMed Abstract](#)
 40. Shimaoka M, Xiao T, Liu JH, *et al.*: Structures of the alpha L I domain and its complex with ICAM-1 reveal a shape-shifting pathway for integrin regulation. *Cell*. 2003; **112**(1): 99–111.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 41. Schubert DA, Gordo S, Sabatino JJ Jr, *et al.*: Self-reactive human CD4 T cell clones form unusual immunological synapses. *J Exp Med*. 2012; **209**(2): 335–352.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 42. Friedman RS, Jacobelli J, Krummel MF: Mechanisms of T cell motility and arrest: deciphering the relationship between intra- and extracellular determinants. *Semin Immunol*. 2005; **17**(6): 387–399.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. Boes M, Cerny J, Massol R, *et al.*: T-cell engagement of dendritic cells rapidly rearranges MHC class II transport. *Nature*. 2002; **418**(6901): 983–988.
[PubMed Abstract](#) | [Publisher Full Text](#)
 44. de la Fuente H, Mittelbrunn M, Sánchez-Martin L, *et al.*: Synaptic clusters of MHC class II molecules induced on DCs by adhesion molecule-mediated initial T-cell scanning. *Mol Biol Cell*. 2005; **16**(7): 3314–3322.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 45. Sakamoto S, Caaveiro JM, Sano E, *et al.*: Contributions of interfacial residues of human Interleukin15 to the specificity and affinity for its private alpha-receptor. *J Mol Biol*. 2009; **389**(5): 880–894.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. Nelson BH, Lord JD, Greenberg PD: Cytoplasmic domains of the interleukin-2 receptor beta and gamma chains mediate the signal for T-cell proliferation. *Nature*. 1994; **369**(6478): 333–336.
[PubMed Abstract](#) | [Publisher Full Text](#)
 47. Mortier E, Woo T, Advincula R, *et al.*: IL-15Ralpha chaperones IL-15 to stable dendritic cell membrane complexes that activate NK cells via trans presentation. *J Exp Med*. 2008; **205**(5): 1213–1225.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 48. Bodnár A, Nizsalóczi E, Mocsár G, *et al.*: A biophysical approach to IL-2 and IL-15 receptor function: localization, conformation and interactions. *Immunol Lett*. 2008; **116**(2): 117–125.
[PubMed Abstract](#) | [Publisher Full Text](#)
 49. Vámosi G, Bodnár A, Vereb G, *et al.*: IL-2 and IL-15 receptor alpha-subunits are coexpressed in a supramolecular receptor cluster in lipid rafts of T cells. *Proc Natl Acad Sci U S A*. 2004; **101**(30): 11082–11087.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 50. Matkó J, Bodnár A, Vereb G, *et al.*: GPI-microdomains (membrane rafts) and signaling of the multi-chain interleukin-2 receptor in human lymphoma/leukemia T cell lines. *Eur J Biochem*. 2002; **269**(4): 1199–1208.
[PubMed Abstract](#) | [Publisher Full Text](#)
 51. Khandelwal S, Roche PA: Distinct MHC class II molecules are associated on the dendritic cell surface in cholesterol-dependent membrane microdomains. *J Biol Chem*. 2010; **285**(46): 35303–35310.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. Chirifu M, Hayashi C, Nakamura T, *et al.*: Crystal structure of the IL-15-IL-15Ralpha complex, a cytokine-receptor unit presented in trans. *Nat Immunol*. 2007; **8**(9): 1001–1007.
[PubMed Abstract](#) | [Publisher Full Text](#)
 53. Stone JD, Conroy LA, Byth KF, *et al.*: Aberrant TCR-mediated signaling in CD45-null thymocytes involves dysfunctional regulation of Lck, Fyn, TCR-zeta, and ZAP-70. *J Immunol*. 1997; **158**(12): 5773–5782.
[PubMed Abstract](#)
 54. Irie-Sasaki J, Sasaki T, Matsumoto W, *et al.*: CD45 is a JAK phosphatase and negatively regulates cytokine receptor signalling. *Nature*. 2001; **409**(6818): 349–354.
[PubMed Abstract](#) | [Publisher Full Text](#)
 55. Varma R, Campi G, Yokosuka T, *et al.*: T cell receptor-proximal signals are sustained in peripheral microclusters and terminated in the central supramolecular activation cluster. *Immunity*. 2006; **25**(1): 117–127.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 56. Dustin ML: Dendritic Cell-Expressed Common Gamma-Chain Recruits IL-15 for Trans-Presentation at the Murine Immunological Synapse. *Open Science Framework*. 2018.
[Data Source](#)

Open Peer Review

Current Referee Status: ? ✓

Version 1

Referee Report 10 September 2018

doi:[10.21956/wellcomeopenres.15778.r33724](https://doi.org/10.21956/wellcomeopenres.15778.r33724)

✓ **Jung-Hyun Park** ¹, **Tae-Hyoun Kim**²

¹ Experimental Immunology Branch, Center for Cancer Research, National Cancer Institute, NIH, Bethesda, MD, USA

² Experimental Immunology Branch, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA

The cytokine IL-15 employs a unique mechanism called “trans-presentation” for signaling. IL-15 transpresentation is distinct to conventional cytokine signaling mechanisms because it requires cell-cell contact between IL-15 expressing and IL-15 signaled cells, and because IL-15 needs to be complexed with its proprietary IL-15Ra-chain to be utilized by target cells. Dendritic cells are potent producers of IL-15. Dendritic cells also play a critical role in antigen presentation to T cells, but it has not been known whether there is crosstalk between these two events. Moreover, it has remained unclear whether these events need to be induced by the same cell or can be triggered by different dendritic cells. Understanding these issues is critical to gain further insights into the regulatory mechanisms of T cell immunity.

The current study by Beilin and colleagues now reports a previously unappreciated requirement for cognate peptide-MHC-II/TCR engagement in IL-15 transpresentation by dendritic cells. Mechanistically, the authors report that the common gamma-chain (gc) cytokine receptor on dendritic cells facilitates IL-15 signaling by recruiting the IL15/IL-15Ra complex to the immunological synapse. Importantly, copatching of gc and IL-15/IL-15Ra was independent of gc receptor signaling, suggesting a recruitment mechanism that is presumably mediated by the receptor ectodomains. Collectively, these findings report a new layer of control in IL-15- transpresentation that directly impacts antigen-specific priming of CD4+ T cells and consequently the establishment of protective T cell immunity.

This study is interesting in two ways:

Firstly, it reveals a previously unappreciated role for gc proteins in IL-15 signaling, that is surprisingly required on IL-15-producing cells – and not on target cells.

Secondly, it reveals a new requirement for TCR ligation to achieve effective IL-15 transpresentation, thus unveiling crosstalk between antigen-presentation and IL-15 transpresentation on the same dendritic cell.

The observations are well documented, and the initial findings are nicely corroborated using a series of imaging studies where cell surface events were scaled down and assessed on lipid bilayers to minimize bystander events. I do not find additional experiments necessary, but I would consider it helpful if the following points could be addressed.

A control experiment that can demonstrate the antigen specificity of the TCR/MHC-II engagement, which

is proposed to be required for IL-15 transpresentation (Figure 1D, E), would be informative. Instead of using OT-II CD4 T cells, can the authors use wildtype CD4+ T cells and co-culture them with OVA preloaded DC? Based on the authors' model, polyclonal wildtype CD4 T cells would fail to phosphorylate STAT5. Is this the case?

Does the recruitment of IL-15/IL-15Ra by gc proteins depend on the cytokine IL-15? Would "empty" IL-15Ra proteins - that are not complexed with IL-15 - suffice to be recruited by gc into the immunological synapse? The authors have established a set of experimental tools in this study that could provide answers to these questions.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

We have read this submission. We believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 25 Sep 2018

Siobhan Burns, Royal Free and University College, UK

Response to comment about antigen specificity:

Using co-cultures between OTII T-cell and wild type DC, we have shown the requirement of MHC-antigen loading for activation of T-cell proliferation and STAT5 phosphorylation (Figure 1 A, D and E) as well as for formation of the DC-T-cell immune synapse in that is a pre-requisite for IL-15 transpresentation in our model (Figure S3). We chose to further dissect whether antigen loading of MHCII is required for T-cell priming using our lipid bilayer model which has the advantage of reducing confounding variables to permit examination of specific elements of antigen presenting cell (APC):T-cell interaction. Data from this approach also supported the need for a cognate interaction between APC and T-cell for IL-15 transpresentation and STAT5 phosphorylation in T-cells (Figure 2A). We agree that future testing of the model we have proposed should include experiments using polyclonal T-cells as suggested to further address the issue of antigen specificity.

Response to comment about 'empty' IL-15Ra:

It was previously shown that IL-15 and IL-15Ra generated by DC form intracellular complexes prior to expression at the cell surface and that these proteins predominantly exist as a complex in DC (Mortier et al JEM 2008, reference 47). Therefore, the degree and physiological relevance of 'empty' IL-15Ra at the immune synapse remains unclear. As a structural question this is interesting as ligation increases the affinity of the interaction of most of the γ c associated alpha chains by 1-2 logs (Gonnord P, Angermann BR, Sadtler K, Gombos E, Chappert P, Meier-Schellersheim M, Varma R. A hierarchy of affinities between cytokine receptors and the common gamma chain leads to pathway cross-talk. Sci Signal. 2018;11(524). doi: 10.1126/scisignal.aal1253. PubMed PMID: 29615515.). Response to point regard Re In these cases the unligated receptor interact with γ c with a 10^{-6} M Kd, which may enable interaction in a synapse as its similar to the affinity of most TCR for agonist pMHC. Therefore, we agree that this is an interesting avenue to explore going forward to test the relative importance of physical interaction and signalling for the model that we have proposed.

Competing Interests: Manuscript author

Referee Report 30 July 2018

doi:[10.21956/wellcomeopenres.15778.r33507](https://doi.org/10.21956/wellcomeopenres.15778.r33507)



Matthew Collin 1,2

¹ Institute of Cellular Medicine, Newcastle University, Newcastle upon Tyne, UK

² NIHR Newcastle BRC, Newcastle Hospitals NHS Foundation Trust, Newcastle, UK

Beilin et al describe in vitro experiments in the murine system that implicate a role for trans-presentation of IL-15 in the activation of CD4+ T cells by GM-CSF stimulated BM-derived dendritic cells. The clinical premise of the work is the observation that γ c and JAK3 SCID patients are not protected from HPV infection after transplantation. The authors surmise that this is due to incomplete chimerism in the myeloid compartment although there are other possibilities as suggested by Reference 2. Laffort et al that uncorrected NK or epithelial cell function could be responsible. The clinical inference made by the authors is tenuous since there are no data on Langerhans cell or dendritic cell chimerism in this patient group as far as I am aware. However, their argument does not detract from the conclusion that IL-15 trans-presentation is shown to be a key player in the immune synapse between MHC II and CD4 T cells.

The authors show that γ c-deficient DC do not have obvious defects of MHCII expression, maturation, antigen uptake and processing, and adhesion to T cells (sup data), yet they are inferior in T cell activation assays (Figure 1). This defect is at least partly reproduced by antibody blockade of IL15R in wild-type cells. The authors go on to dissect the mechanism of IL-15 trans-presentation showing that T cell STAT5 is only activated in the context of TCR ligation (Figure 2) that γ c and IL-15R are recruited to the synapse (Figure 3,4,5) and that signalling to the DC is not required, through the use of a γ c C-terminal truncation (Figure 5).

Points requiring clarification

1) The GM-CSF stimulated BM DC has recently come under intense scrutiny as a heterogeneous preparation of DC-lineage and monocyte-lineage cells (e.g Helft, J¹). The explanation that IL15 trans-presentation is the major defect in γ c -/- preparations critically rests on these preparations being functionally equivalent (in all other respects) to WT cells. Given that GM-CSF also signals through

STAT5, it is important to exclude an interaction between GM-CSF and γc in the generation of $\gamma c^{-/-}$ DC. For example, γc cytokines originating from a plethora of cells in the BM could enhance the function of wild-type preparations in some way. The authors could improve their description of BM-derived DC by presenting more detailed flow cytometry or bulk gene expression analysis of wild-type and $\gamma c^{-/-}$ preparations. The methods section should at least contain an outline of how the cells were obtained (the use of GM-CSF was gleaned from the results sections).

2) In Figure 1, it would be useful to have more clarity concerning how much the defect in antigen presentation by $\gamma c^{-/-}$ DC was attributable to IL-15, e.g in panel F the $\gamma c^{-/-}$ are normalised to their own poor performance. Why not show the raw data rather than % inhibition? Related to this point, the authors cite a notable previous study in which trans-presentation of IL-2 was demonstrated to be essential at the immune synapse (14. Wuest et al). It was not clear from the data or interpretation what the relative importance of IL15 and IL2 trans-presentation would be since both could contribute to the inferior performance observed in the $\gamma c^{-/-}$ phenotype. This point could receive more attention in the discussion.

References

1. Helft J, Böttcher J, Chakravarty P, Zelenay S, Huotari J, Schraml BU, Goubau D, Reis e Sousa C: GM-CSF Mouse Bone Marrow Cultures Comprise a Heterogeneous Population of CD11c(+)MHCII(+) Macrophages and Dendritic Cells. *Immunity*. 2015; **42** (6): 1197-211 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Referee Expertise: Human dendritic cell and macrophage ontogeny and function

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 25 Sep 2018

Siobhan Burns, Royal Free and University College, UK

Response to Point 1:

We agree with the reviewer that bone marrow-derived dendritic cells (BMDC), while widely adopted, represent an imperfect model of in vivo counterparts. The reviewer makes an important point about whether absence of γc may alter the differentiation of bone marrow cells. In the original manuscript we present evidence that γc -deficient BMDC generated using the protocol described are phenotypically comparable with wild type (WT) controls with respect to MHCII expression and upregulation of the maturation marker CD86 and that several dendritic cell-associated functions, such as antigen uptake and presentation, are the same between γc -deficient and WT BMDC. We have additional data to show that γc -deficient and WT BMDC are not different in expression of the DC surface marker CD11c or the maturation marker CD80 (new Figure S2C,E). We did not specifically analyse CD11b subsets within our cultures (as done in the Helft et al manuscript highlighted by the reviewer) to define BMDC from macrophages but we did not see any significant differences in expression of CD80 and CD86 in CD11c+ cells which would be predicted if the proportions of BMDC and macrophages were altered (as these were specifically identified as markers poorly expressed in bone marrow derived macrophages). Furthermore, we did not see any significant differences in release of the proinflammatory cytokines IL-6 or TNF- α following stimulation with TLR ligands including CpG (which was also shown by Helft et al to differ between BMDC and macrophages; new Figure S2K). Thus, while we cannot exclude minor differences, our data indicate that the phenotype of cells generated from γc -deficient bone marrow using GM-CSF are broadly comparable with WT control.

In support of our in vitro differentiation data, we also observed that the ex-vivo phenotype of splenic DC was comparable between γc -deficient and WT mice (Figure S1 A,B,C). While there were significantly fewer CD11c+ DCs in the spleens of γc -deficient mice when compared to WT controls ($p < 0.05$), as previously described for other lymphopenic mouse models (Asli B, Lantz O, DiSanto JP, Saeland S, Geissmann F. Roles of lymphoid cells in the differentiation of Langerhans dendritic cells in mice. *Immunobiology*. 2004;209(1-2):209-21; PMID: 15481155), within the CD11c-enriched fraction, WT and γc /Rag 2 $^{-/-}$ mice had similar frequencies of CD11c+ CD11b+ and CD11c+ CD8 α + conventional DCs. CD11c+ B220+ plasmacytoid DC were also present in comparable numbers in the spleens of WT and γc /Rag 2 $^{-/-}$ mice. Thus, these results suggest that although required for DC development in vivo, γc is probably dispensable for DC differentiation.

For this study, bone-marrow (BM) cells were extracted from the femur and tibia of mice. BM-derived DCs were grown from BM cells cultured over 7 days in RPMI medium 1640 supplemented with 10% fetal bovine serum (FBS) and 1% Penicillin/Streptomycin (Gibco) in the presence of 20 ng/ml GM-CSF (BioSource). For all experiments, BM-derived DCs were CD11c selected using magnetic bead separation (Miltenyi Biotech). To induce DC activation, CD11c+ DCs were matured overnight with 100ng/ml LPS (Sigma). This detail has been added to the methods section for clarity.

Response to Point 2:

Raw data from a representative example is shown in the original manuscript in supplemental data (now Figure S4C). Addition of IL-15Ra blocking antibody reduced the ability of WT BMDC to induce T-cell proliferation, as assessed by flow cytometry measurement of CFSE dilution, from 54% to 37% at maximal antibody dose (10mcg/ml). The level of proliferation induced by WT BMDC in the presence of IL-15 blockade was similar to the level of T-cell proliferation induced by

$\gamma c^{-/-}$ DC without blockade (37%), suggesting that the defect of T-cell priming seen with $\gamma c^{-/-}$ DC was almost entirely attributable to defective IL-15 transpresentation. Furthermore, there was no real change in T-cell proliferation when $\gamma c^{-/-}$ DC were co-cultured with anti-IL15Ra blocking antibody (reduced from 37% to 36%) indicating that the effect seen in WT cells was specific. With respect to the relative importance of IL-15 and IL-2 trans presentation, we tested this experimentally in BMDC: T-cell coculture assays in which we reasoned that BMDC would be transpresenting IL-15 or IL-2 to induce phosphorylation of STAT5 in T-cells. We observed complete abrogation of T-cell pSTAT if we added anti-IL15Ra blocking antibody to the co-culture (Figure 1D) but no change in T-cell pSTAT5 using anti-IL2Ra blockade (Figure S4A,B). This led us to conclude that 'STAT5 activation occurs primarily through IL-15 transpresentation during DC-mediated priming of naïve CD4+ T cells in our experimental system'. We have added the following sentence to the discussion (end of paragraph2):

'Although it has been previously been shown that blockade of the DC IL-2Ra reduces T-cell activation (Wuest et al, ref 14), we were unable to demonstrate a contribution of DC-mediated IL-2Ra transpresentation in our system leading us to conclude that IL-15 is the major cytokine transpresented by DC for T-cell priming.'

Competing Interests: Manuscript author